



# Longitudinal Changes in Objectively-Measured Physical Activity and Sedentary Time among School-Age Children in Central Texas, US during the COVID-19 Pandemic

Leigh Ann Ganzar, PhD<sup>1</sup>; Deborah Salvo, PhD<sup>2</sup>; Katie Burford, MS<sup>1</sup>; Yuzi Zhang, MS<sup>1</sup>; Harold W. Kohl III, PhD<sup>1,2</sup>; Deanna M. Hoelscher, PhD<sup>1</sup>

<sup>1</sup>Michael & Susan Dell Center for Healthy Living, UTHealth School of Public Health in Austin, Texas, USA; <sup>2</sup>Department of Kinesiology and Health Education, The University of Texas at Austin, Texas, USA



## INTRODUCTION

- Regular physical activity is important for energy-balance in children.
- COVID-19 affected how people live, work, study, travel, and play.
- Previous evidence on physical activity during COVID-19:
  - Cross-sectional studies
  - Self-report measures
  - Outside of US

## AIMS

1. To identify change trajectories of device-measured physical activity and sedentary time from pre-COVID-19 to during COVID-19 in school-aged children in the US
2. To examine the socio-ecological factors associated with changes in movement behaviors.

## METHODS

- **Design:**
  - Part of STREETS 5-year natural experiment
  - Longitudinal study design with two time points:
    - **Time 1:** Sept 2019-Feb 2020 (Pre-COVID-19)
    - **Time 2:** October 2020-March 2021 (During COVID-19)
- **Population Sample:** Cohort of school-age children (ages 8-11)
- **Measures:**
  - GT3X accelerometers using Evenson cut points to measure primary outcomes of:
    - Mean daily minutes of moderate-to-vigorous physical activity (MVPA)
    - Mean daily hours sedentary time
  - Socio-ecological predictors: individual, family, social, organizational, neighborhood
- **Data analysis methods**
  - Descriptive statistics
  - Latent class linear mixed models (Aim 1)
  - Logistic regression models (Aim 2)

## ACKNOWLEDGEMENTS & REFERENCES

### References

1. Yang S, Guo B, Ao L, Yang C, Zhang L, Zhou J, et al. Obesity and activity patterns before and during COVID-19 lockdown among youths in China. *Clinical Obesity*. 2020;10(6):e12416.
2. Yamada K, Kurita S. Influence of social distancing during the COVID-19 pandemic on physical activity in children: A scoping review of the literature. *Journal of Exercise Science & Fitness*. 2021.
3. Okely AD, Karipipattan KE, Guan H, Taylor EK, Suesse T, Cross PL, et al. Global effect of COVID-19 pandemic on physical activity, sedentary behaviour and sleep among 3-to 5-year-old children: a longitudinal study of 14 countries. *BMC Public Health*. 2021;21(1):1-15.
4. Burkart S, Parker H, Weaver RG, Beets MW, Jones A, Adams EL, et al. Impact of the COVID-19 pandemic on elementary schoolers' physical activity, sleep, screen time and diet: A quasi-experimental interrupted time series study. *Pediatric Obesity*. 2022;17(1):e12846
5. Cooper AR, Goodman A, Page AS, Sherar LB, Esliger DW, van Sluijs EM, et al. Objectively measured physical activity and sedentary time in youth: the international children's accelerometry database (ICAD). *Int J Behav Nutr Phys Act*. 2015; 12(1): 113

### Acknowledgements

The STREETS Study would like to thank the City of Austin Public Works Department, and all participating school districts, campuses, and participants for their investment in the study.

The STREETS study is funded by the Eunice Kennedy Shriver National Institute of Child Health & Human Development, grant number R01 HD097669 and the Michael & Susan Dell Foundation through the Michael & Susan Dell Center for Healthy Living.

## RESULTS

**168** Number of participants with valid physical activity at both timepoints

**56% female**  
**44% male**

**9** Average age at baseline in years

**44% White, Non-Hispanic**  
**39% Hispanic or Latinx**  
**10% Asian or Other**  
**7% Black or African American**


**29%** with parents who have high school education or less

**54%** virtual school attendance during COVID



Scan for more results!

Figure 1. Physical Activity Trajectories



### Important factors:

- Gender: girls less likely to maintain MVPA
- Social cohesion: higher cohesion more likely to maintain MVPA


### Average daily MVPA changes:

- Decrease MVPA group: -11.10 minutes
- Maintain High MVPA: -1.66 minutes

### Latent Class

- 1: 'Decrease MVPA', n=138
- 2: 'Maintain High MVPA', n=30

Figure 2. Sedentary Behavior Trajectories



### Important factors:

- Race/ethnicity: Hispanic children more likely to decrease ST
- Social cohesion: higher cohesion less likely to decrease ST

### Average daily ST changes:

- Moderate increase ST: 0.95 hours
- Steep increase ST: 3.40 hours
- Decrease ST: -0.75 hours

### Latent Class

- 1: 'Moderate Increase Sedentary', n=132
- 2: 'Steep Increase Sedentary', n=10
- 3: 'Decrease Sedentary', n=26

## CONCLUSIONS & NEXT STEPS

- Significant declines in physical activity and increases in sedentary time during the COVID-19 pandemic when compared to pre-COVID-19 time period.
- The majority of children in this study were categorized in the 'decreasing MVPA' and 'increasing sedentary time' groups.
- Previous evidence yearly relative change in minutes of daily MVPA from age 3 to 18 was -3.4%. We found a mean yearly relative change in minutes of daily MVPA of -17.0%.

### Limitations

- Small sample size
- Maturation bias
- Determinants of movement behaviors not measured

### Future research

Despite these limitations, this study demonstrates the need to counteract short-term negative changes to children's movement behaviors in the face of societal level disruptors.